



Ozone-cci+



Product Validation Plan (PVP)

Date: 28/11/2025

Version: 4.1

Phase 3, Task 2

Deliverable D2.4

WP Manager: V. Sofieva

WP Manager Organization: FMI

Other partners: AUTH, BIRA-IASB



DOCUMENT PROPERTIES

Title Product Validation Plan (PVP)
Reference Ozone_cci+_D2.4_PVP_v4.1
Issue / Revision 4 / 1
Status Final
Date of issue 28/11/2025
Document type Validation Protocol

	FUNCTION	NAME	DATE
LEAD AUTHORS	VALT leader	Arno Keppens, BIRA-IASB	11/09/2025
CONTRIBUTING AUTHORS	VALT members	Jean-Christopher Lambert, BIRA-IASB Tijl Verhoelst, BIRA-IASB José Granville, BIRA-IASB Katerina Garane, AUTH Dimitris Balis, AUTH	11/09/2025
REVIEWED BY	Science leader	Daan Hubert, BIRA-IASB	13/10/2025
ACCEPTED BY	ESA Officer	Michael Eisinger, ESA	28/11/2025
ISSUED BY	Science leader	Daan Hubert, BIRA-IASB	28/11/2025



DOCUMENT CHANGE RECORD

Issue	Revision	Date	Modified items	Observations
1	0	03/01/2012	First version.	
2	0	20/01/2020	Completely revised version to align with CCI+ objectives and with state-of-the-art developments.	
2	1	06/12/2020	Updated applicable documents (Section 2) and references to the updated URD v3.1 (Section 3).	
3	0	30/09/2023	Completely revised version to align with CCI+ Phase 2 objectives and with state-of-the-art developments.	
3	1	17/03/2024	Minor revisions upon ESA feedback.	
4	0	26/06/2025	Revised version to align with CCI+ Phase 3 objectives and developments.	
4	1	28/11/2025	Minor revisions and removal of Section 3 (User Requirements) following ESA feedback.	



Table of Contents

1	Introduction.....	7
1.1	Purpose and scope	7
1.2	Document overview	7
2	Applicable and reference documents	8
2.1	Requirement documents	8
2.2	Standards and framework documents	8
2.3	Validation references.....	9
2.3.1	Ozone column validation	9
2.3.2	Nadir ozone profile validation	11
2.3.3	Limb ozone profile validation.....	12
2.3.4	Tropospheric ozone column validation.....	15
2.3.5	Methods, merged data products validation and miscellaneous.....	16
4	ECV Product Evaluation Protocol	19
4.1	Foreword	19
4.2	Generic principles applicable to all ECV products	19
4.2.1	Core requirements of the GEOSS data quality strategy (QA4EO).....	19
4.2.2	Principles of the validation of atmospheric data.....	19
4.2.3	Principles of the validation of an ECV product line	19
4.2.4	Confrontation with independent reference data.....	21
4.2.4.1	Generalities	21
4.2.4.2	Reference measurements from GAW ground-based networks	21
4.2.4.3	Error budget of a data comparison	21
4.2.4.4	Information content	22
4.2.5	Validation of individual components.....	23
4.2.6	Validation against service specifications	23
4.2.7	Validation against user requirements.....	23
4.2.8	Quality control of operational ECV data production	23
4.2.9	Validation of ECV product updates	23
4.3	Validation specifics by ECV product.....	24
4.3.1	Total ozone data product.....	24
4.3.1.1	Validation requirements	24
4.3.1.2	Validation data sources.....	24
4.3.2	Ozone profile data product from nadir-viewing instruments.....	25
4.3.2.1	Validation requirements	25
4.3.2.2	Validation data sources.....	25
4.3.3	Ozone profile data product from limb-viewing instruments.....	26
4.3.3.1	Validation requirements	26
4.3.3.2	Validation data sources.....	26
4.3.4	Tropospheric ozone data products	27
4.3.4.1	Validation requirements	27
4.3.4.2	Validation data sources.....	27
5	Standards.....	28
5.1	Maintenance of datasets and reports	28
5.2	Metadata and additional information	28
5.3	QA and validation metadata.....	28
5.4	Compliance with international standards	29



6	Terms and definitions	31
6.1	Terms and definitions.....	31
6.2	Abbreviations and acronyms	34



1 Introduction

1.1 Purpose and scope

This Product Validation Plan (PVP) summarises the validation requirements for the ozone Essential Climate Variable (ECV) data products of ESA's Ozone_cci+ project, namely, the total ozone column, the tropospheric ozone column and vertical ozone profiles based on nadir and on limb/occultation satellite measurements. This plan updates earlier versions developed and applied in the CCI programme, aligning with the objectives of Phase 3 of the Ozone_cci+ project.

1.2 Document overview

This Ozone_cci+ Product Validation Plan is organised as follows:

- Chapter 1 contains this introduction describing the scope of the document.
- Chapter 2 lists applicable and reference documents.
- Chapter 3 defines the Evaluation Protocol for the final ECV product. It starts with generic principles of the ECV validation and explains the specifics with regard to validation of the different ozone ECVs.
- Chapter 4 addresses validation and quality control standards: sustainable archiving and traceability of the validation process and of validation results, quality control metadata and criteria, and compliance with international standards.
- Chapter 6 defines the recommended terminology, abbreviations and acronyms.



2 Applicable and reference documents

2.1 Requirement documents

- [RD1] CMUG: Climate Community Requirements, Deliverable 1.1, Climate Modelling User Group, version 4, May 2024.
- [RD2] DARD: Ozone CCI Data Access Requirement Document, version 2.1, Ozone_cci_DARD_2.1, 25/05/2016.
- [RD3] WMO/GCOS: World Meteorological Organization (WMO); Intergovernmental Oceanographic Commission of the United Nations Educational, Scientific and Cultural Organization (IOC-UNESCO); International Science Council (ISC); United Nations Environment Programme (UNEP); Copernicus Climate Change Service (C3S), The 2022 GCOS ECVs Requirements 2022 edition - Updated in 2025, GCOS-245, <https://library.wmo.int/idurl/4/58111>.
- [RD4] IGACO: The changing atmosphere. An integrated global atmospheric chemistry observation theme for the IGOS partnership. Report of the Integrated Global Atmospheric Chemistry Observation (IGACO) theme team, September 2004 (ESA SP-1282, GAW No. 159, WMO-TD No. 1235), 2004.
- [RD5] URD: Ozone CCI User Requirements Document, Version 5.1, Ozone_cci+_D1.1_URD_v5.1, 28/11/2025.
- [RD6] WMO: OSCAR (Observing Systems Capability Analysis and Review Tool, <https://www.wmo-sat.info/oscar/observingrequirements> (last access June, 2025).

2.2 Standards and framework documents

- [RD7] CDRH: Center for Devices and Radiological Health (CDRH), General Principles of Software Validation; Final Guidance for Industry and FDA Staff, CBER CDRH/OC #938, 11/01/2002. Publicly available via <http://www.fda.gov/MedicalDevices/DeviceRegulationandGuidance>
- [RD8] CEOS: Committee on Earth Observation Satellites (CEOS): Terms and Definitions and other documents and resources, publicly available on <http://calvalportal.ceos.org> (last access June, 2025).
- [RD9] GUM: Joint Committee for Guides in Metrology (JCGM/WG 1) 100:2008, Evaluation of measurement data – Guide to the expression of uncertainty in a measurement (GUM), http://www.bipm.org/utils/common/documents/jcgm/JCGM_100_2008_E.pdf
- [RD10] Larssen, S., R. Sluyter, and C. Helmis, Criteria for EUROAIRNET – The EEA Air Quality Monitoring and Information Network, <https://www.eea.europa.eu/publications/TEC12>, 1999.
- [RD11] Nappo, C.J., Caneill J.Y., Furman R.W., Gifford F.A., Kaimal J.C., Kramer M.L., Lockhart T.J., Pendergast M.M., Pielke R.A., Randerson D., Shreffler J.H., and Wyngaard J.C., The Workshop on the Representativeness of Meteorological Observations, June 1981, Boulder, CO, Bull. Am. Meteorol. Soc. 63, 761-764, <http://www.jstor.org/stable/26222836>, 1982.
- [RD12] NIST: Prokhorov, A. V., R. U. Datla, V. P. Zakharenkov, V. Privalsky, T. W. Humpherys, and V. I. Sapritsky, Spaceborne Optoelectronic Sensors and their Radiometric Calibration. Terms and Definitions. Part 1. Calibration Techniques, Ed. by A. C. Parr and L. K. Issaev, NIST Technical Note NISTIR 7203, March 2005.
- [RD13] VIM: Joint Committee for Guides in Metrology (JCGM/WG 2) 200:2008 & ISO/IEC Guide 99-12:2007, International Vocabulary of Metrology – Basic and General Concepts and Associated Terms (VIM), <http://www.bipm.org/en/publications/guides/vim.html>



[RD14] WMO Quality Management Framework (QMF), home page at <https://public.wmo.int/en/our-mandate/how-we-do-it/quality-management-framework>

[RD15] QA4EO – A Quality Assurance framework for Earth Observation, established by the CEOS. It consists of ten distinct key guidelines linked through an overarching document (the QA4EO Guidelines Framework) and more community-specific QA4EO procedures, all available on <http://qa4eo.org/documentation>. A short QA4EO "user" guide has been produced to provide background into QA4EO and how one would start implementing it (http://qa4eo.org/docs/QA4EO_guide.pdf)

[RD16] ISO Quality Management Principles available at <https://asq.org/quality-resources/iso-9000> and <https://asq.org/quality-resources/iso-14000>

[RD17] NetCDF Climate and Forecast Metadata Convention, <http://cfconventions.org>

[RD18] Fahre Vik, A., T. Krognes, S-E. Walker, S. Bjørndalsæter, C. Stoll, T. Bårde, R. Paltiel, and B. Gloslie, ESA Campaign Database (CDB) user manual, NILU Technical Note O-103045, 100 pp., April 2006. <https://www.nilu.no/dnn/ACF830.pdf>

[RD19] World Meteorological Organization, Global Atmosphere Watch (GAW) Implementation Plan: 2016-2023, GAW Report No. 228, https://library.wmo.int/doc_num.php?explnum_id=3395.

2.3 Validation references

2.3.1 Ozone column validation

[RD20] Antón, M., M. E. Koukouli, M. Kroon, R. D. McPeters, G. J. Labow, D. Balis, and A. Serrano, Global validation of empirically corrected EP Total Ozone Mapping Spectrometer (TOMS) total ozone columns using Brewer and Dobson ground-based measurements, *J. Geophys. Res.*, 115, D19305, doi:10.1029/2010JD014178, 2010.

[RD21] Balis, D., J-C. Lambert, M. Van Roozendael, D. Loyola, R. Spurr, Y. Livschitz, P. Valks, V. Amiridis, P. Gerard, and J. Granville, Ten years of GOME/ERS-2 total ozone data – The new GOME Data Processor (GDP) Version 4: II Ground-based validation and comparisons with TOMS V7/V8, *J. Geophys. Res.*, Vol. 112, D07307, doi:10.1029/2005JD006376, 2007.

[RD22] Balis, D., M. Kroon, M. E. Koukouli, E. J. Brinksma, G. Labow, J. P. Veefkind, and R. D. McPeters, Validation of Ozone Monitoring Instrument total ozone column measurements using Brewer and Dobson spectrophotometer ground-based observations, *J. Geophys. Res.*, 112, D24S46, doi:10.1029/2007JD008796, 2007.

[RD23] Bracher, A., Lamsal, L. N., Weber, M., Bramstedt, K., Coldewey-Egbers, M., and Burrows, J. P., Global satellite validation of SCIAMACHY O₃ columns with GOME WFDOAS, *Atmos. Chem. Phys.*, 5, 2357-2368, doi:10.5194/acp-5-2357-2005, 2005.

[RD24] Bramstedt, K., J. Gleason, D. Loyola, W. Thomas, A. Bracher, M. Weber, and J. P. Burrows, Comparison of total ozone from the satellite instruments GOME and TOMS with measurements from the Dobson network 1996–2000, *Atmos. Chem. Phys.*, 3, 1409-1419, doi:10.5194/acp-3-1409-2003, 2003.

[RD25] Coldewey-Egbers, M., Loyola, D. G., Koukouli, M., Balis, D., Lambert, J.-C., Verhoelst, T., Granville, J., van Roozendael, M., Lerot, C., Spurr, R., Frith, S. M., and Zehner, C.: The GOME-type Total Ozone Essential Climate Variable (GTO-ECV) data record from the ESA Climate Change Initiative, *Atmos. Meas. Tech.*, 8, 3923-3940, doi:10.5194/amt-8-3923-2015, 2015.

[RD26] Fioletov, V. E., G. Labow, R. Evans, *et al.*, Performance of the ground-based total ozone network assessed using satellite data, *J. Geophys. Res.*, Vol. 113, D14313, doi:10.1029/2008JD009809, 2008.



[RD27] Garane, K., Lerot, C., Coldewey-Egbers, M., Verhoelst, T., Koukouli, M. E., Zyrichidou, I., Balis, D. S., Danckaert, T., Goutail, F., Granville, J., Hubert, D., Keppens, A., Lambert, J.-C., Loyola, D., Pommereau, J.-P., Van Roozendael, M., and Zehner, C.: Quality assessment of the Ozone_cci Climate Research Data Package (release 2017) – Part 1: Ground-based validation of total ozone column data products, *Atmos. Meas. Tech.*, 11, 1385-1402, <https://doi.org/10.5194/amt-11-1385-2018>, 2018.

[RD28] Koukouli, M., D. Balis, I. Zyrichidou, C. Lerot, M. Van Roozendael, J-C. Lambert, J. Granville, J-P. Pommereau, F. Goutail, G. Labow, S. Frith, D. Loyola, R. Spurr, and C. Zehner, Evaluating a new homogeneous total ozone climate data record from GOME/ERS-2, SCIAMACHY/Envisat, and GOME-2/MetOp-A, *J. Geophys. Res. Atmos.*, 120(23), 12,212–296,312, doi:10.1002/2015JD023699, 2015.

[RD29] Koukouli, M. E., Zara, M., Lerot, C., Frakos, K., Balis, D., van Roozendael, M., Allart, M. A. F., and van der A, R. J.: The impact of the ozone effective temperature on satellite validation using the Dobson spectrophotometer network, *Atmos. Meas. Tech.*, 9, 2055-2065, doi:10.5194/amt-9-2055-2016, 2016.

[RD30] Lambert, J.-C., D. S. Balis, P. Gerard, J. Granville, Y. Livschitz, D. Loyola, R. Spurr, P. Valks, and M. Van Roozendael, UPAS / GDOAS 4.0 Upgrade of the GOME Data Processor for Improved Total Ozone Columns – Delta Validation Report, Ed. by J.-C. Lambert (IASB) and D. Balis (AUTH), Tech. Note ERSE-CLVL-EOPG-TN-04-0001, European Space Agency, Frascati, Italy, 2004.

[RD31] Lambert, J.-C., G. Hansen, V. Soebijanta, W. Thomas, M. Van Roozendael, D. S. Balis, C. Fayt, P. Gerard, J. F. Gleason, J. Granville, G. Labow, D. Loyola, J. H. G. van Geffen, R. F. van Oss, C. Zehner, and C. S. Zerefos,, ERS-2 GOME GDP3.0 implementation and validation, Edited by J.-C. Lambert (IASB), Tech. Note ERSE-DTEXEOAD-TN-02-0006, 138 pp., European Space Agency, Frascati, Italy, 2002.

[RD32] Lambert, J.-C., M. E. Koukouli, D. S. Balis, J. Granville, C. Lerot, D. Pieroux, and M. Van Roozendael, GDP 5.0 - Upgrade of the GOME Data Processor for Improved Total Ozone Column - Validation Report for ERS-2 GOME GDP 5.0 Total Ozone Column, Edited by J.-C. Lambert (IASB) and M. E. Koukouli (AUTH), Tech. Note TN-IASB-GOME-GDP5-VR, Issue/Rev. 1/A, 55 pp., 6 May 2011.

[RD33] Lambert, J.-C., M. Van Roozendael, M. De Mazière, P.C. Simon, J.-P. Pommereau, F. Goutail, A. Sarkissian, and J.F. Gleason, Investigation of pole-to-pole performances of spaceborne atmospheric chemistry sensors with the NDSC, *J. Atmos. Sci.*, Vol. 56, 176-193, doi: 10.1175/1520-0469, 1999.

[RD34] Lambert, J.-C., M. Van Roozendael, P.C. Simon, J.-P. Pommereau, F. Goutail, J.F. Gleason, S.B. Andersen, D.W. Arlander, N.A. Bui Van, H. Claude, J. de La Noë, M. De Mazière, V. Dorokhov, P. Eriksen, A. Green, K. Karlsen Tørnkvist, B.A. Kåstad Høiskar, E. Kyrö, J. Leveau, M.-F. Merienne, G. Milinevsky, H.K. Roscoe, A. Sarkissian, J.D. Shanklin, J. Staehelin, C. Wahlstrøm Tellefsen, and G. Vaughan, Combined characterisation of GOME and TOMS total ozone measurements from space using ground-based observations from the NDSC, *Adv. Space Res.*, Vol. 26, 1931-1940, 2000.

[RD35] Loyola, D. G., M. E. Koukouli, P. Valks, D. S. Balis, N. Hao, M. Van Roozendael, R. J. D. Spurr, W. Zimmer, S. Kiemle, C. Lerot, and J-C. Lambert, The GOME-2 Total Column Ozone Product: Retrieval Algorithm and Ground-Based Validation, *J. Geophys. Res.*, Vol. 116, doi:10.1029/2010JD014675, 2011.



[RD36] Verhoelst, T., Granville, J., Hendrick, F., Köhler, U., Lerot, C., Pommereau, J.-P., Redondas, A., Van Roozendael, M., and Lambert, J.-C.: Metrology of ground-based satellite validation: co-location mismatch and smoothing issues of total ozone comparisons, *Atmos. Meas. Tech.*, 8, 5039-5062, doi:10.5194/amt-8-5039-2015, 2015.

[RD37] Weber, M., Lamsal, L. N., Coldewey-Egbers, M., Bramstedt, K., and Burrows, J. P., Pole-to-pole validation of GOME WFDOAS total ozone with groundbased data, *Atmos. Chem. Phys.*, 5, 1341-1355, doi:10.5194/acp-5-1341-2005, 2005.

2.3.2 Nadir ozone profile validation

[RD38] Bracher, A., M. Weber, K. Bramstedt, S. Tellmann, and J. P. Burrows, Long-term global measurements of ozone profiles by GOME validated with SAGE II considering atmospheric dynamics, *J. Geophys. Res.*, 109, D20308, doi:10.1029/2004JD004677, 2004.

[RD39] De Clercq, C., J.-C. Lambert, O. Tuinder, and R. van Oss, Tropospheric ozone information in GOME long-term data record, in *Proc. Envisat Symposium 2007, Montreux, Switzerland, 23-27 April 2007*, ESA Special Publication SP-636, 7 pp., 2007.

[RD40] De Clercq, C., J.-C. Lambert, J. Granville, P. Gerard, A. Kaifel, J. Kaptur, B. Mijling, O. tuinder, R. van Oss, and C. Zehner, CHEOPS-GOME, Geophysical information content and validation of ERS-2 GOME ozone profile data records, IASB/ESA Technical Note TN-IASB-GOME1-CHEOPS-01, Issue 1, Revision B, 122 pp., 20 December 2007.

[RD41] Keppens, A., Lambert, J.-C., Granville, J., Miles, G., Siddans, R., van Peet, J. C. A., van der A, R. J., Hubert, D., Verhoelst, T., Delcloo, A., Godin-Beekmann, S., Kivi, R., Stübi, R., and Zehner, C.: Round-robin evaluation of nadir ozone profile retrievals: methodology and application to MetOp-A GOME-2, *Atmos. Meas. Tech.*, 8, 2093-2120, doi:10.5194/amt-8-2093-2015, 2015.

[RD42] Keppens, A., Lambert, J.-C., Granville, J., Hubert, D., Verhoelst, T., Compernolle, S., Latter, B., Kerridge, B., Siddans, R., Boynard, A., Hadji-Lazaro, J., Clerbaux, C., Wespes, C., Hurtmans, D. R., Coheur, P.-F., van Peet, J. C. A., van der A, R. J., Garane, K., Koukouli, M. E., Balis, D. S., Delcloo, A., Kivi, R., Stübi, R., Godin-Beekmann, S., Van Roozendael, M., and Zehner, C.: Quality assessment of the Ozone_cci Climate Research Data Package (release 2017) – Part 2: Ground-based validation of nadir ozone profile data products, *Atmos. Meas. Tech.*, 11, 3769-3800, <https://doi.org/10.5194/amt-11-3769-2018>, 2018.

[RD43] Keppens, A., Compernolle, S., Verhoelst, T., Hubert, D., and Lambert, J.-C.: Harmonization and comparison of vertically resolved atmospheric state observations: methods, effects, and uncertainty budget, *Atmos. Meas. Tech.*, 12, 4379–4391, <https://doi.org/10.5194/amt-12-4379-2019>, 2019.

[RD44] Keppens, A.; Di Pede, S.; Hubert, D.; Lambert, J.-C.; Veefkind, P.; Sneep, M.; De Haan, J.; ter Linden, M.; Leblanc, T.; Compernolle, S.; Verhoelst, T.; Granville, J.; Nath, O.; Fjaeraa, A. M.; Boyd, I.; Niemeijer, S.; Van Malderen, R.; Smit, H. G. J.; Duflat, V.; Godin-Beekmann, S.; Johnson, B. J.; Steinbrecht, W.; Tarasick, D. W.; Kollonige, D. E.; Stauffer, R. M.; Thompson, A. M.; Dehn, A. & Zehner, C. 5 years of Sentinel-5P TROPOMI operational ozone profiling and geophysical validation using ozonesonde and lidar ground-based networks *Atmospheric Measurement Techniques*, 2024, 17, 3969-3993, doi:10.5194/amt-17-3969-2024.

[RD45] Liu, X., K. Chance, C. E. Sioris, T. P. Kurosu, and M. J. Newchurch, Intercomparison of GOME, ozonesonde, and SAGE II measurements of ozone: Demonstration of the need to homogenize available ozonesonde data sets, *J. Geophys. Res.*, 111, D14305, doi:10.1029/2005JD006718, 2006.



[RD46] Liu, X., K. Chance, C. E. Sioris, R. J. D. Spurr, T. P. Kurosu, R. V. Martin, and M. J. Newchurch, Ozone profile and tropospheric ozone retrievals from the Global Ozone Monitoring Experiment: Algorithm description and validation, *J. Geophys. Res.*, 110, D20307, doi:10.1029/2005JD006240, 2005.

[RD47] Liu, X., K. Chance, and T. P. Kurosu, Improved ozone profile retrievals from GOME data with degradation correction in reflectance, *Atmos. Chem. Phys.*, 7, 1575-1583, doi:10.5194/acp-7-1575-2007, 2007.

[RD48] Meijer, Y. J., D. P. J. Swart, F. Baier, P. K. Bhartia, G. E. Bodeker, S. Casadio, K. Chance, F. Del Frate, T. Erbertseder, L. E. Flynn, S. Godin-Beekmann, G. Hansen, O. P. Hasekamp, A. Kaifel, H. M. Kelder, B. J. Kerridge, J.-C. Lambert, J. Landgraf, B. Latter, X. Liu, I. S. McDermid, M. D. Müller, Y. Pachepsky, V. Rozanov, R. Siddans, S. Tellmann, R. J. van der A, R. F. van Oss, M. Weber, and C. Zehner, Evaluation of Global Ozone Monitoring Experiment (GOME) ozone profiles from nine different algorithms, *J. Geophys. Res.*, Vol. 111, D21306, doi:10.1029/2005JD006778, 2006.

[RD49] Miles, G. M., Siddans, R., Kerridge, B. J., Latter, B. G., and Richards, N. A. D.: Tropospheric ozone and ozone profiles retrieved from GOME-2 and their validation, *Atmos. Meas. Tech.*, 8, 385-398, doi:10.5194/amt-8-385-2015, 2015.

[RD50] Nassar, R., Logan, J. A., Worden, H. M., Megretskaya, I. A., Bowman, K. W., Osterman, G. B., Thompson, A. M., Tarasick, D. W., Austin, S., Claude, H., Dubey, M. K., Hocking, W. K., Johnson, B. J., Joseph, E., Merrill, J., Morris, G. A., Newchurch, M., Oltmans, S. J., Posny, F., Schmidlin, F. J., Vomel, H., Whiteman, D. N., and Witte, J. C., Validation of Tropospheric Emission Spectrometer (TES) nadir ozone profiles using ozonesonde measurements, *J. Geophys. Res.*, 113, D15S17, doi:10.1029/2007JD008819, 2008.

[RD51] Osterman, G. B., S. S. Kulawik, H. M. Worden, N. A. D. Richards, B. M. Fisher, A. Eldering, M. W. Shephard, L. Froidevaux, G. Labow, M. Luo, R. L. Herman, K. W. Bowman, and A. M. Thompson, Validation of Tropospheric Emission Spectrometer (TES) measurements of the total, stratospheric, and tropospheric column abundance of ozone, *J. Geophys. Res.*, 113, D15S16, doi:10.1029/2007JD008801, 2008.

[RD52] Timmermans, R. M. A., R. F. van Oss, and H. M. Kelder, Equatorial Kelvin wave signatures in ozone profile measurements from Global Ozone Monitoring Experiment (GOME), *J. Geophys. Res.*, 110, D21103, doi:10.1029/2005JD005929, 2005.

2.3.3 Limb ozone profile validation

[RD53] Adams, C., A. E. Bourassa, A. F. Bathgate, C. A. McLinden, N. D. Lloyd, C. Z. Roth, E. J. Llewellyn, J. M. Zawodny, D. E. Flittner, G. L. Manney, W. H. Daffer, and D. A. Degenstein, Characterization of Odin-OSIRIS ozone profiles with the SAGE II dataset, *Atmos. Meas. Tech.*, 6, 1447-1459, 2013.

[RD54] Adams, C., A. E. Bourassa, V. Sofieva, L. Froidevaux, C. A. McLinden, D. Hubert, J. -C. Lambert, C. E. Sioris, and D. A. Degenstein, Assessment of Odin-OSIRIS ozone measurements from 2001 to the present using MLS, GOMOS, and ozone sondes, *Atmos. Meas. Tech.*, 7, 49-64, doi:10.5194/amt-7-49-2014, 2014.

[RD55] Brinksma, E., A. Bracher, D. E. Lolkema, A. J. Segers, I. S. Boyd, K. Bramstedt, H. Claude, S. Godin-Beekmann, G. Hansen, G. Kopp, T. Leblanc, I. S. McDermid, Y. J. Meijer, H. Nakane, A. Parrish, C. von Savigny, K. Stebel, D. P. J. Swart, G. Taha, and A. J. M. Piters, Geophysical validation of SCIAMACHY Limb Ozone Profiles, *Atmos. Chem. Phys.*, 6, 197-209, doi:10.5194/acp-6-197-2006, 2006.



[RD56] Cortesi, U., J.-C. Lambert, C. De Clercq, G. Bianchini, T. Blumenstock, A. Bracher, E. Castelli, V. Catoire, K. V. Chance, M. De Mazière, P. Demoulin, S. Godin-Beekmann, N. Jones, K. Jucks, C. Keim, T. Kerzenmacher, H. Kuellmann, J. Kuttippurath, M. Iarlori, G. Y. Liu, Y. Liu, I. S. McDermid, Y. J. Meijer, F. Mencaraglia, S. Mikuteit, H. Oelhaf, C. Piccolo, M. Pirre, P. Raspollini, F. Ravegnani, W. J. Reburn, G. Redaelli, J. J. Remedios, H. Sembhi, D. Smale, T. Steck, A. Taddei, C. Varotsos, C. Vigouroux, A. Waterfall, G. Wetzel, and S. Wood, Geophysical validation of MIPAS-Envisat operational ozone data, *Atmos. Chem. Phys.*, Vol. 7, 4807-4867, 2007.

[RD57] Dupuy, E., J. Kar, K. Walker, P. Bernath, G. Bodeker, C. Boone, I. Boyd, A. Bracher, V. Catoire, T. Christensen, U. Cortesi, J. Davies, C. De Clercq, L. Froidevaux, D. Fussen, P. von der Gathen, F. Goutail, C. Haley, L. Harvey, R. Hughes, J. Jin, A. Jones, A. Kagawa, Y. Kasai, T. Kerzenmacher, A. Klekociuk, J.-C. Lambert, N. Lloyd, E. Mahieu, G. Manney, C.T. McElroy, S. McLeod, A. Parrish, S. Petelina, C. Piccolo, C. Randall, C. Roth, C. von Savigny, T. Steck, K. Strong, R. Sussmann, A. Thompson, M. Tully, and J. Urban, Validation of ozone measurements from the Atmospheric Chemistry Experiment, *Atmos. Chem. Phys.*, Vol. 9, 287-343, 2009.

[RD58] Froidevaux, L., Y. B. Jiang, A. Lambert, N. J. Livesey, W. G. Read, J. W. Waters, E. V. Browell, J. W. Hair, M. A. Avery, T. J. McGee, L. W. Twigg, G. K. Sumnicht, K. W. Jucks, J. J. Margitan, B. Sen, R. A. Stachnik, G. C. Toon, P. F. Bernath, C. D. Boone, K. A. Walker, M. J. Filipiak, R. S. Harwood, R. A. Fuller, G. L. Manney, M. J. Schwartz, W. H. Daffer, B. J. Drouin, R. E. Cofield, D. T. Cuddy, R. F. Jarnot, B. W. Knosp, V. S. Perun, W. V. Snyder, P. C. Stek, R. P. Thurstans, and P. A. Wagner, Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *J. Geophys. Res.*, 113, D15S20, doi:10.1029/2007JD008771, 2008.

[RD59] Hubert, D., Lambert, J.-C., Verhoelst, T., Granville, J., Keppens, A., Baray, J.-L., Bourassa, A. E., Cortesi, U., Degenstein, D. A., Froidevaux, L., Godin-Beekmann, S., Hoppel, K. W., Johnson, B. J., Kyrölä, E., Leblanc, T., Lichtenberg, G., Marchand, M., McElroy, C. T., Murtagh, D., Nakane, H., Portafaix, T., Querel, R., Russell III, J. M., Salvador, J., Smit, H. G. J., Stebel, K., Steinbrecht, W., Strawbridge, K. B., Stübi, R., Swart, D. P. J., Taha, G., Tarasick, D. W., Thompson, A. M., Urban, J., van Gijsel, J. A. E., Van Malderen, R., von der Gathen, P., Walker, K. A., Wolfram, E., and Zawodny, J. M.: Ground-based assessment of the bias and long-term stability of 14 limb and occultation ozone profile data records, *Atmos. Meas. Tech.*, 9, 2497-2534, doi:10.5194/amt-9-2497-2016, 2016.

[RD60] Jiang, Y. B., L. Froidevaux, A. Lambert, N. J. Livesey, W. G. Read, J. W. Waters, B. Bojkov, T. Leblanc, I. S. McDermid, S. Godin-Beekmann, M. J. Filipiak, R. S. Harwood, R. A. Fuller, W. H. Daffer, B. J. Drouin, R. E. Cofield, D. T. Cuddy, R. F. Jarnot, B. W. Knosp, V. S. Perun, M. J. Schwartz, W. V. Snyder, P. C. Stek, R. P. Thurstans, P. A. Wagner, M. Allaart, S. B. Andersen, G. Bodeker, B. Calpini, H. Claude, G. Coetzee, J. Davies, H. De Backer, H. Dier, M. Fujiwara, B. Johnson, H. Kelder, N. P. Leme, G. König-Langlo, E. Kyro, G. Laneve, L. S. Fook, J. Merrill, G. Morris, M. Newchurch, S. Oltmans, M. C. Parrondos, F. Posny, F. Schmidlin, P. Skrivankova, R. Stubi, D. Tarasick, A. Thompson, V. Thouret, P. Viatte, H. Vömel, P. von Der Gathen, M. Yela, and G. Zablocki, Validation of Aura Microwave Limb Sounder Ozone by ozonesonde and lidar measurements, *J. Geophys. Res.*, 112, D24S34, doi:10.1029/2007JD008776, 2007.

[RD61] Keckhut, P., Hauchecorne, A., Blanot, L., Hocke, K., Godin-Beekmann, S., Bertaux, J.-L., Barrot, G., Kyrölä, E., van Gijsel, J. A. E., and Pazmino, A.: Mid-latitude ozone monitoring with the GOMOS-ENVISAT experiment version 5: the noise issue, *Atmos. Chem. Phys.*, 10, 11839-11849, doi:10.5194/acp-10-11839-2010, 2010.



[RD62] Laeng, A., Grabowski, U., von Clarmann, T., Stiller, G., Glatthor, N., Höpfner, M., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., Sofieva, V., Petropavlovskikh, I., Hubert, D., Bathgate, T., Bernath, P., Boone, C. D., Clerbaux, C., Coheur, P., Damadeo, R., Degenstein, D., Frith, S., Froidevaux, L., Gille, J., Hoppel, K., McHugh, M., Kasai, Y., Lumpe, J., Rahpoe, N., Toon, G., Sano, T., Suzuki, M., Tamminen, J., Urban, J., Walker, K., Weber, M., and Zawodny, J., Validation of MIPAS IMK/IAA V5R_O3_224 ozone profiles, *Atmos. Meas. Tech.*, 7, 3971-3987, doi:10.5194/amt-7-3971-2014, 2014.

[RD63] Laeng, A.; Hubert, D.; Verhoelst, T.; von Clarmann, T.; Dinelli, B.M.; Dudhia, A.; Raspollini, P.; Stiller, G.; Grabowski, U.; Keppens, A.; Kiefer, M. ; Sofieva, V.; Froidevaux, L.; Walker, K.; Lambert, J.-C.; Zehner, C., The ozone climate change initiative: Comparison of four Level-2 processors for the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS), *Remote Sens. Environ.*, doi:<http://dx.doi.org/10.1016/j.rse.2014.12.013>, 2015.

[RD64] Manney, G. L., W. H. Daffer, J. M. Zawodny, P. F. Bernath, K. W. Hoppel, K. A. Walker, B. W. Knosp, C. Boone, E. E. Remsberg, M. L. Santee, V. L. Harvey, S. Pawson, D. R. Jackson, L. Deaver, C. T. McElroy, C. A. McLinden, J. R. Drummond, H. C. Pumphrey, A. Lambert, M. J. Schwartz, L. Froidevaux, S. McLeod, L. L. Takacs, M. J. Suarez, C. R. Trepte, D. C. Cuddy, N. J. Livesey, R. S. Harwood, and J. W. Waters, Solar occultation satellite data and derived meteorological products: Sampling issues and comparisons with Aura Microwave Limb Sounder, *J. Geophys. Res.*, 112, D24S50, doi:10.1029/2007JD008709, 2007.

[RD65] Meijer, Y. J., D. P. J. Swart, R. Koelemeijer, M. Allaart, S. Andersen, G. Bodeker, I. Boyd, G. Braathen, Y. Calisesi, H. Claude, V. Dorokhov, P. von der Gathen, M. Gil, S. Godin-Beekmann, F. Goutail, G. Hansen, A. Karpetchko, P. Keckhut, H. Kelder, B. Kois, R. Koopman, J.-C. Lambert, T. Leblanc, I. S. McDermid, S. Pal, U. Raffalski, H. Schets, R. Stubi, T. Suortti, G. Visconti, and M. Yela, Pole-to-pole validation of ENVISAT GOMOS ozone profiles using data from ground-based and balloon-sonde measurements, *J. Geophys. Res.*, Vol. 109, D23305, doi:10.1029/2004JD004834, 2004.

[RD66] Mieruch, S., M. Weber, C. von Savigny, A. Rozanov, H. Bovensmann, J. P. Burrows, P. F. Bernath, C. D. Boone, L. Froidevaux, L. L. Gordley, M. G. Mlynczak, J. M. Russell III, L. W. Thomason, K. A. Walker, and J. M. Zawodny, Global and long-term comparison of SCIAMACHY limb ozone profiles with correlative satellite data (2002-2008), *Atmos. Meas. Tech.*, 5, 771-788, doi:10.5194/amt-5-771-2012, 2012.

[RD67] Mze, N., Hauchecorne, A., Bencherif, H., Dalaudier, F., and Bertaux, J.-L.: Climatology and comparison of ozone from ENVISAT/GOMOS and SHADOZ/balloon-sonde observations in the southern tropics, *Atmos. Chem. Phys.*, 10, 8025-8035, doi:10.5194/acp-10-8025-2010, 2010.

[RD68] Rahpoe, N., von Savigny, C., Weber, M., Rozanov, A.V., Bovensmann, H., and Burrows, J. P.: Error budget analysis of SCIAMACHY limb ozone profile retrievals using the SCIATRAN model, *Atmos. Meas. Tech.*, 6, 2825-2837, doi:10.5194/amt-6-2825-2013, 2013.

[RD69] Rahpoe, N., Weber, M., Rozanov, A. V., Weigel, K., Bovensmann, H., Burrows, J. P., Laeng, A., Stiller, G., von Clarmann, T., Kyrölä, E., Sofieva, V. F., Tamminen, J., Walker, K., Degenstein, D., Bourassa, A. E., Hargreaves, R., Bernath, P., Urban, J., and Murtagh, D. P., Relative drifts and biases between six ozone limb satellite measurements from the last decade, *Atmos. Meas. Tech.*, 8, 4369-4381, doi:10.5194/amt-8-4369-2015, 2015.

[RD70] Rohen, G. J., C. von Savigny, J. W. Kaiser, E. J. Llewellyn, L. Froidevaux, M. López-Puertas, T. Steck, M. Palm, H. Winkler, M. Sinnhuber, H. Bovensmann, and J. P. Burrows, Ozone profile retrieval from limb scatter measurements in the HARTLEY bands: further retrieval details and profile comparisons, *Atmos. Chem. Phys.*, 8, doi:10.5194/acp-8-2509-2008, 2509-2517, 2008.



[RD71] Sofieva, V. F., Rahpoe, N., Tamminen, J., Kyrölä, E., Kalakoski, N., Weber, M., Rozanov, A., von Savigny, C., Laeng, A., von Clarmann, T., Stiller, G., Lossow, S., Degenstein, D., Bourassa, A., Adams, C., Roth, C., Lloyd, N., Bernath, P., Hargreaves, R. J., Urban, J., Murtagh, D., Hauchecorne, A., Dalaudier, F., van Roozendael, M., Kalb, N., and Zehner, C.: Harmonized dataset of ozone profiles from satellite limb and occultation measurements, *Earth Syst. Sci. Data*, 5, 349-363, doi:10.5194/essd-5-349-2013, 2013.

[RD72] Sofieva, V. F., Tamminen, J., Kyrölä, E., Laeng, A., von Clarmann, T., Dalaudier, F., Hauchecorne, A., Bertaux, J.-L., Barrot, G., Blanot, L., Fussen, D. and Vanhellemont, F.: Validation of GOMOS ozone precision estimates in the stratosphere, *Atmos. Meas. Tech.*, 7(7), 2147-2158, doi:10.5194/amt-7-2147-2014, 2014.

[RD73] Van Gijsel, J. A. E., Swart, D. P. J., Baray, J.-L., Bencherif, H., Claude, H., Fehr, T., Godin-Beekmann, S., Hansen, G. H., Keckhut, P., Leblanc, T., McDermid, I. S., Meijer, Y. J., Nakane, H., Quel, E. J., Stebel, K., Steinbrecht, W., Strawbridge, K. B., Tatarov, B. I., and Wolfram, E. A.: GOMOS ozone profile validation using ground-based and balloon sonde measurements, *Atmos. Chem. Phys.*, 10, 10473-10488, doi:10.5194/acp-10-10473-2010, 2010.

2.3.4 Tropospheric ozone column validation

[RD74] Arosio, C., Sofieva, V., Orfanoz-Cheuquela, A., Rozanov, A., Heue, K.-P., Malina, E., Stauffer, R. M., Tarasick, D., Van Malderen, R., Ziemke, J. R., and Weber, M.: Inter-comparison of tropospheric ozone column datasets from combined nadir and limb satellite observations, EGUSphere [preprint], <https://doi.org/10.5194/egusphere-2024-3737>, in review, 2024.

[RD75] Cooper, O. R.; Parrish, D. D.; Ziemke, J.; Balashov, N. V.; Cupeiro, M.; Galbally, I. E.; Gilge, S.; Horowitz, L.; Jensen, N. R.; Lamarque, J.-F. & et al. Global distribution and trends of tropospheric ozone: An observation-based review *Elem. Sci. Anth.*, BioOne, 2014, 2, 000029.

[RD76] Ebojie, F.; von Savigny, C.; Ladstätter-Weißenmayer, A.; Rozanov, A.; Weber, M.; Eichmann, K.; Bötel, S.; Rahpoe, N.; Bovensmann, H. & Burrows, J. P. Tropospheric column amount of ozone retrieved from SCIAMACHY limb-nadir-matching observations *Atmos. Meas. Tech.*, 2014, 6, 7811-7865.

[RD77] Heue, K.-P.; Coldewey-Egbers, M.; Delcloo, A.; Lerot, C.; Loyola, D.; Valks, P. & van Roozendael, M. Trends of tropical tropospheric ozone from 20 years of European satellite measurements and perspectives for the Sentinel-5 Precursor *Atmos. Meas. Tech.*, 2016, 9, 5037-5051.

[RD78] Hubert, D.; Heue, K.-P.; Lambert, J.-C.; Verhoelst, T.; Allaart, M.; Compernolle, S.; Cullis, P. D.; Dehn, A.; Félix, C.; Johnson, B. J.; Keppens, A.; Kollonige, D. E.; Lerot, C.; Loyola, D.; Maata, M.; Mitro, S.; Mohamad, M.; Piters, A.; Romahn, F.; Selkirk, H. B.; da Silva, F. R.; Stauffer, R. M.; Thompson, A. M.; Veefkind, J. P.; Vömel, H.; Witte, J. C. & Zehner, C. TROPOMI tropospheric ozone column data: geophysical assessment and comparison to ozonesondes, GOME-2B and OMI *Atmos. Meas. Tech.*, 2021, 14, 7405-7433.

[RD79] Keim, C.; Eremenko, M.; Orphal, J.; Dufour, G.; Flaud, J.-M.; Höpfner, M.; Boynard, A.; Clerbaux, C.; Payan, S.; Coheur, P.-F.; Hurtmans, D.; Claude, H.; Dier, H.; Johnson, B.; Kelder, H.; Kivi, R.; Koide, T.; López Bartolomé, M.; Lambkin, K.; Moore, D.; Schmidlin, F. J. & Stübi, R. Tropospheric ozone from IASI: comparison of different inversion algorithms and validation with ozone sondes in the northern middle latitudes *Atmos. Chem. Phys.*, 2009, 9, 9329-9347.

[RD80] Miles, G. M.; Siddans, R.; Kerridge, B. J.; Latter, B. G. & Richards, N. A. D. Tropospheric ozone and ozone profiles retrieved from GOME-2 and their validation *Atmos. Meas. Tech.*, 2015, 8, 385-398.



[RD81] Pope, R. J., O'Connor, F. M., Dalvi, M., Kerridge, B. J., Siddans, R., Latter, B. G., Barret, B., Le Flochmoen, E., Boynard, A., Chipperfield, M. P., Feng, W., Pimplott, M. A., Dhomse, S. S., Retscher, C., Wespes, C., and Rigby, R.: Investigation of the impact of satellite vertical sensitivity on long-term retrieved lower-tropospheric ozone trends, *Atmos. Chem. Phys.*, 24, 9177–9195, <https://doi.org/10.5194/acp-24-9177-2024>, 2024.

[RD82] Zhang, L. & others Intercomparison methods for satellite measurements of atmospheric composition: application to tropospheric ozone from TES and OMI *Atmos. Chem. Phys.*, 2010, 10, 4725-4739.

[RD83] Ziemke, J. R.; Chandra, S.; Duncan, B. N.; Froidevaux, L.; Bhartia, P. K.; Levelt, P. F. & Waters, J. W. Tropospheric ozone determined from Aura OMI and MLS: Evaluation of measurements and comparison with the Global Modeling Initiative's Chemical Transport Model *J. Geophys. Res.*, 2006, 111.

2.3.5 Methods, merged data products validation and miscellaneous

[RD84] Calisesi, Y., V. T. Soebijanta, and R. van Oss, Regridding of remote soundings: Formulation and application to ozone profile comparison, *J. Geophys. Res.*, 110, D23306, doi:10.1029/2005JD006122, 2005.

[RD85] Chiou, E. W., Bhartia, P. K., McPeters, R. D., Loyola, D. G., Coldewey-Egbers, M., Fioletov, V. E., Van Roozendael, M., Spurr, R., Lerot, C., and Frith, S. M.: Comparison of profile total ozone from SBUV (v8.6) with GOME-type and ground-based total ozone for a 16-year period (1996 to 2011), *Atmos. Meas. Tech.*, 7, 1681-1692, doi:10.5194/amt-7-1681-2014, 2014.

[RD86] Coldewey-Egbers, M., Loyola R., D. G., Braesicke, P., Dameris, M., Van Roozendael, M., Lerot, C., and Zimmer, W., A new health check of the ozone layer at global and regional scales. *Geophysical Research Letters*, Vol.41, pp. 4363-4372, doi:10.1002/2014GL060212, 2014.

[RD87] De Clercq, C., J.-C. Lambert, J. Granville, P. Gerard, A. Kaifel, J. Kaptur, and C. Zehner, CHEOPS-GOME: Soundness of new ozone profile climatologies, IASB/ESA Technical Note TN-IASB-GOME1-CHEOPS-02, Issue 1, Revision B, 29 pp., 27 June 2007.

[RD88] Dirksen, R. J., Sommer, M., Immler, F. J., Hurst, D. F., Kivi, R., and Vömel, H.: Reference quality upper-air measurements: GRUAN data processing for the Vaisala RS92 radiosonde, *Atmos. Meas. Tech.*, 7, 4463–4490, <https://doi.org/10.5194/amt-7-4463-2014>, 2014.

[RD89] Eckert, E., T. von Clarmann, M. Kiefer, G. P. Stiller, S. Lossow, N. Glatthor, D. A. Degenstein, L. Froidevaux, S. Godin-Beekmann, T. Leblanc, S. McDermid, M. Pastel, W. Steinbrecht, D. P. J. Swart, K. A. Walker, and P. F. Bernath, Drift-corrected trends and periodic variations in MIPAS IMK/IAA ozone measurements, *Atmos. Chem. Phys.*, 14, 2571-2589, doi:10.5194/acp-14-2571-2014, 2014.

[RD90] Keppens, A., Compernolle, S., Hubert, D., Verhoelst, T., Granville, J., and Lambert, J.-C.: Removing Prior Information from Remotely Sensed Atmospheric Profiles by Wiener Deconvolution Based on the Complete Data Fusion Framework, *Remote Sens.*, 14, 2197, <https://doi.org/10.3390/rs14092197>, 2022.

[RD91] Errera, Q., F. Daerden, S. Chabriat, J.-C. Lambert, W. A. Lahoz, S. Viscardy, S. Bonjean, and D. Fonteyn, 4D-Var Assimilation of MIPAS Chemical Observations: Ozone and Nitrogen Dioxide Analyses, *Atmos. Chem. Phys.*, Vol. 8, 6169-6187, 2008.

[RD92] Kyrölä, E., M. Laine, V. Sofieva, J. Tamminen, S.-M. Päivärinta, S. Tukiainen, J. Zawodny, and L. Thomason, Combined SAGE II-GOMOS ozone profile data set 1984–2011 and trend analysis of the vertical distribution of ozone, *Atmos. Chem. Phys.*, 13, 10645-10658, doi:10.5194/acp-13-10645-2013, 2013.



[RD93] Lambert, J.-C., De Clercq C., von Clarmann T., Combining and Merging Water Vapour Observations: A Multi-dimensional Perspective on Smoothing and Sampling Issues. In: Kämpfer N. (eds) Monitoring Atmospheric Water Vapour. ISSI Scientific Report Series, vol 10. Springer, New York, NY, 2013.

[RD94] Lauer, A., V. Eyring, M. Righi, M. Buchwitz, P. Defourny, M. Evaldsson, P. Friedlingstein, R. de Jeu, G. de Leeuw, A. Loew, C. J. Merchant, B. Müller, T. Popp, M. Reuter, S. Sandven, D. Senftleben, M. Stengel, M. Van Roozendael, S. Wenzel, U. Willén, Benchmarking CMIP5 models with a subset of ESA CCI Phase 2 data using the ESMValTool, *Remote Sens. Environ.*, doi:<http://dx.doi.org/10.1016/j.rse.2017.01.007>, 2017.

[RD95] Lerot, C., M. Van Roozendael, R. Spurr, D. Loyola, M. Coldewey-Egbers, S. Kochanova, J. van Gent, M. Koukouli, D. Balis, J.-C. Lambert, J. Granville, C. Zehner, Homogenized total ozone data records from the European sensors GOME/ERS-2, SCIAMACHY/Envisat and GOME-2/Metop-A, *J. Geophys. Res.*, 119, 1–20, doi:10.1002/2013JD020831, 2014.

[RD96] Loew, A., Bell, W., Brocca, L., Bulgin, C.E., Burdanowitz, J., Calbet, X., Donner, R.V., Ghent, D., Gruber, A., Kaminski, T., Kinzel, J., Klepp, C., Lambert, J.-C., Schaepman-Strub, G., Schröder, M., Verhoelst, T., Validation practices for satellite-based Earth observation data across communities, *Rev. Geophys.*, 55, 779– 817, doi:10.1002/2017RG000562, 2017.

[RD97] Meijer, Y. J., R. J. van der A, R. F. van Oss, D. P. J. Swart, H. M. Kelder, and P. V. Johnston, Global Ozone Monitoring Experiment ozone profile characterization using interpretation tools and lidar measurements for intercomparison, *J. Geophys. Res.*, Vol. 108 (D23), 4723, doi:10.1029/2003JD003498, 2003.

[RD98] Merchant, C. J., Paul, F., Popp, T., Ablain, M., Bontemps, S., Defourny, P., Hollmann, R., Lavergne, T., Laeng, A., de Leeuw, G., Mittaz, J., Poulsen, C., Povey, A. C., Reuter, M., Sathyendranath, S., Sandven, S., Sofieva, V. F., and Wagner, W.: Uncertainty information in climate data records from Earth observation, *Earth Syst. Sci. Data*, 9, 511-527, <https://doi.org/10.5194/essd-9-511-2017>, 2017.

[RD99] Müller, M. D., A. K. Kaifel, M. Weber, S. Tellmann, J. P. Burrows, and D. Loyola, Ozone profile retrieval from Global Ozone Monitoring Experiment (GOME) data using a neural network approach (Neural Network Ozone Retrieval System (NNORSY)), *J. Geophys. Res.*, 108(D16), 4497, doi:10.1029/2002JD002784, 2003.

[RD100] Nair, P. J., S. Godin-Beekmann, A. Pazmiño, A. Hauchecorne, G. Ancellet, I. Petropavlovskikh, L. E. Flynn, and L. Froidevaux, Coherence of long-term stratospheric ozone vertical distribution time series used for the study of ozone recovery at a northern mid-latitude station, *Atmos. Chem. Phys.*, 11, 4957-4975, doi:10.5194/acp-11-4957-2011, 2011.

[RD101] Rodgers, C. D., and B. J. Connor, Intercomparison of remote sounding instruments, *J. Geophys. Res.*, 108, 4116, doi:10.1029/2002JD002299, 2003.

[RD102] Sofieva, V. F., Kalakoski, N., Päivärinta, S.-M., Tamminen, J., Laine, M., and Froidevaux, L.: On sampling uncertainty of satellite ozone profile measurements, *Atmos. Meas. Tech.*, 7, 1891– 1900, doi:10.5194/amt-7-1891-2014, 2014.

[RD103] Sofieva, V. F., Kyrölä, E., Laine, M., Tamminen, J., Degenstein, D., Bourassa, A., Roth, C., Zawada, D., Weber, M., Rozanov, A., Rahpoe, N., Stiller, G., Laeng, A., von Clarmann, T., Walker, K. A., Sheese, P., Hubert, D., van Roozendael, M., Zehner, C., Damadeo, R., Zawodny, J., Kramarova, N., and Bhartia, P. K.: Merged SAGE II, Ozone_cci and OMPS ozone profile dataset and evaluation of ozone trends in the stratosphere, *Atmos. Chem. Phys.*, 17, 12533-12552, <https://doi.org/10.5194/acp-17-12533-2017>, 2017.



[RD104] van Peet, J. C. A., van der A, R. J., Tuinder, O. N. E., Wolfram, E., Salvador, J., Levelt, P. F., and Kelder, H. M.: Ozone Profile Retrieval Algorithm (OPERA) for nadir-looking satellite instruments in the UV–VIS, *Atmos. Meas. Tech.*, 7, 859-876, doi:10.5194/amt-7-859-2014, 2014.

[RD105] van Peet, J. C. A., van der A, R. J., Kelder, H. M., and Levelt, P. F.: Simultaneous assimilation of ozone profiles from multiple UV-VIS satellite instruments, *Atmos. Chem. Phys.*, 18, 1685-1704, <https://doi.org/10.5194/acp-18-1685-2018>, 2018.

[RD106] Viscardy, S., Q. Errera, Q., Y. Christophe, S. Chabriat, and J.-C. Lambert, Evaluation of ozone analyses from UARS MLS assimilation by BASCOE between 1992 and 1997, *Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, Vol. 3:2, 190-202, doi: 10.1109/JSTARS.2010.2040463, 2010.

[RD107] von Clarmann, T., Validation of remotely sensed profiles of atmospheric state variables: strategies and terminology, *Atmos. Chem. Phys.*, 6, 4311–4320, doi:10.5194/acp-6-4311-2006, 2006 + Addendum: *Atmos. Chem. Phys.*, 6, 5547-5547, 2006.

[RD108] von Clarmann, T., and U. Grabowski, Elimination of hidden a priori information from remotely sensed profile data, *Atmos. Chem. Phys.*, 7, 397-408, doi:10.5194/acp-7-397-2007, 2007.

[RD109] von Clarmann, T., C. De Clercq, M. Ridolfi, M. Höpfner, and J.-C. Lambert, The horizontal resolution of MIPAS, *Atmos. Meas. Tech.*, Vol. 2, 47-54, 2009.

[RD110] von Clarmann, T., G. Stiller, U. Grabowski, E. Eckert, and J. Orphal, Technical Note: Trend estimation from irregularly sampled, correlated data, *Atmos. Chem. Phys.*, 10, 6737-6747, doi:10.5194/acp-10-6737-2010, 2010.

[RD111] Weber, M., Coldewey-Egbers, M., Fioletov, V. E., Frith, S. M., Wild, J. D., Burrows, J. P., Long, C. S., and Loyola, D.: Total ozone trends from 1979 to 2016 derived from five merged observational datasets – the emergence into ozone recovery, *Atmos. Chem. Phys.*, 18, 2097-2117, <https://doi.org/10.5194/acp-18-2097-2018>, 2018.



4 ECV Product Evaluation Protocol

4.1 Foreword

This chapter starts with the general principles applicable to the validation of the ozone ECV products. It continues with the specific characteristics applicable for each of the ozone ECV products. As a baseline, generic principles and means for validation shall prevail over specific provisions whenever possible, in order to enable a standardised approach. This chapter applies to the full validation of the final ECV products.

4.2 Generic principles applicable to all ECV products

4.2.1 Core requirements of the GEOSS data quality strategy (QA4EO)

The Quality Assurance Framework for Earth Observation (QA4EO) [RD15] establishes general principles of the data quality strategy for the Global Earth Observation System of Systems (GEOSS). The core requirement of QA4EO is that all data and derived products shall have associated with them a documented and fully traceable quality indicator (QI). This quality indicator shall provide sufficient information to allow all users to readily evaluate the “fitness for purpose” of the data or derived product. The quality indicator shall be based on a documented and quantifiable assessment of evidence demonstrating the level of traceability to internationally agreed (where possible, SI) reference standards.

4.2.2 Principles of the validation of atmospheric data

The validation of an atmospheric ozone data product can be seen as a science-driven verification process, the aim of which being to verify that the data produced do respond to predefined quality requirements and information content requirements. Validation generally involves the assessment of the closeness of the data to the geophysical reality, and of its sources of uncertainty, over the spatial and temporal domains of relevance as defined in the URD. Uncertainty estimates can include, but are not restricted to, estimates of the bias and dispersion of the data with respect to reference data, and identification of the temporal and spatial domains over which those estimates are valid. Standard concepts of the classical metrology, like precision and repeatability, usually apply to atmospheric measurements. However, they can be of limited suitability for modelling results, for which more dedicated quality indicators shall be defined. It must be noted that international standardisation bodies insist on the fact that accuracy – defined as the closeness of agreement between a quantity value obtained by measurement and the true value of the measurand – is not a quantity and hence is not given a numerical quantity value [RD13].

4.2.3 Principles of the validation of an ECV product line

In a metrology-like approach of validation, the quality of data products must be evaluated (1) through assessment of uncertainties associated with the way the data product is measured or calculated, and (2) through confrontation with ‘reference’ measurements showing documented evidence of quality traceable to international standards, following community agreed practices [RD96]. In the context of CCI, quality must be evaluated also through critical analysis of the suitability of the data products for the targeted applications, i.e. through the validation of the actual usability of the datasets.

Figure 1 presents an overview of the main validation tasks and quality control mechanisms to be applied over the life cycle of every ECV data production.




Figure 1. Validation tasks, quality control mechanisms and feedback loops over the life cycle of an ozone ECV data production line.

From top to bottom, the box chart shows the timeline for the evolution of an ECV data production chain (centre column blue square boxes) through phases from the build-up through operations to updates and its associated validation steps (right-hand column of orange square boxes). The high-level appointment of responsibilities is outlined in the centre column (oval boxes), highlighting the respective role of research partners, of system developers and ECV data producers (ALGT), of validation teams (VALT) and climate research users (CRG), and of ECV data producers in the general QA/QC loop. Major feedback loops are also highlighted, from those associated to operations feedback into improvements of algorithms and their operationalisation into ECV data production lines, to the formal endorsement by CRG users. The latter step concludes officially the build-up of an operational service. The following sections describe the major validation tasks in more detail.



4.2.4 Confrontation with independent reference data

4.2.4.1 Generalities

The performance of calibration procedures, retrieval algorithms and merging systems, and the quality of the resulting ECV products will be assessed by comparison with reference measurements providing the atmospheric “truth”. A key aspect of any comparison for validation purposes is the selection of the reference data sets. The quality, traceability and suitability of the latter are essential to allow proper, unbiased and independent validation. Reference measurements must be well documented and procedures must exist to ensure adequate quality control on the long term, as it is the case, e.g., within international ground-based networks.

Where and when reference observations are available, they constitute the preferred source of validation data, superseding the use of modelling results as validation data. When suitable measurements are not available, validation of data might also involve comparisons with “reference” model data sets. Models are of valuable use to extend measurement-based validation to the global domain and to a better sampling of temporal and spatial features, to verify data products under atmospheric states and scenarios not accessible to the measurement, to assess comparison errors due to temporal and spatial mismatch and differences in sampling, and to identify inconsistencies in the data sets under investigation. However models, including data assimilation systems, must always be used with circumspection in validation as they are based on our current understanding of the atmosphere and our current ability to model/algorithmically depict this understanding and they can suffer from many limitations and uncertainties.

4.2.4.2 Reference measurements from GAW ground-based networks

Ground-based reference measurements of the total column and vertical distribution of ozone are performed by networks of instruments contributing to WMO’s Global Atmosphere Watch programme (GAW) [RD19]. Data sets suitable for the validation analysis of ECV products are collected from complementary instruments archiving routinely their data to the World Ozone and Ultraviolet Radiation Data Centre (WOUDC) and the Data Host Facility (DHF) of the Network for the Detection of Atmospheric Composition Change (NDACC). Individual details are given in the Data Access Requirement Document (DARD) [RD2]. Access conditions and pricing as applicable to the two data archives are regulated by data protocols available on the web portals of the data archives (<http://woudc.org> and <http://ndacc.org>, respectively).

It should be pointed out that the Ozone CCI project does not foresee the production of any independent validation measurements. It needs to rely completely on observations and results provided by existing monitoring networks and ongoing/planned research projects as described in the DARD. High-level impetus through ESA, CEOS, the EC, space agencies and national agencies funding instrument operation as part of networks, is also required, in particular to ensure data provision suitable for sustainable validation activities of the future operational ECV data production.

4.2.4.3 Error budget of a data comparison

A major objective of quantitative comparisons with reference measurements is to estimate the validity of the theoretical (ex-ante) uncertainties provided with the data product. However, the discrepancy between the satellite data set being validated and the reference data set combines uncertainties associated with each individual system, plus uncertainties associated with the methodology of comparison. Discrepancies include the effect of the following comparison uncertainties:

- (1) Comparison uncertainties associated with the difference in sampling of atmospheric variability and structures: e.g. geographical mismatch, diurnal cycle effects in the upper stratosphere and mesosphere (USM), assumptions related to the area of representativeness.



(2) Comparison uncertainties associated with the difference in smoothing of atmospheric variability and structures: e.g., balloon-based in situ measurement at about 150 m resolution by an electrochemical cell, compared with GOME ground pixels of $40 \times 320 \text{ km}^2$ or TROPOMI ground pixels of $5.5 \times 3.5 \text{ km}^2$ and a vertical resolution of 3-8 km.

As much as possible, most comparison uncertainties will be reduced by a cautious design of the selection of data sets to be compared, and by considering that a multivariate analysis of the comparison results taking into account the specifics of the data being compared (modelling data or remote sensing data, atmospheric variability and gradients etc.) might be required and preferred over entirely statistical approaches. For traceability purposes it is essential to document, for each validation exercise, the selection method applied to the data sets (temporal and spatial co-location criteria, how differences in vertical and horizontal smoothing are handled etc.) [RD43].

Although essential, the derivation of a complete error budget for each comparison is still a matter of research at the time being [RD36] and it falls partly beyond the scope of the Ozone_cci+ project. Validation teams (VALT) as well as data producers (ALGT) are aware that neglecting uncertainties linked to the comparison method can spoil the value of the comparison and yield erroneous conclusions on the quality of the compared data product. This awareness must be transmitted to the reader of Ozone_cci+ Validation Reports for a proper use of the validation results and, in fine, of the ozone ECV data records. When misinterpretation is possible, common statements like “the discrepancy between the two data sets ranges within their individual error bars” will be suitably annexed with a provision on the – actual calculated or simply expected – contribution of the selection and comparison methods to this discrepancy. Provisions like “temporal and spatial mismatches exist but their contribution to the discrepancy between the two data sets has not been assessed; nevertheless this contribution is assumed to be small...” or “the selection method has been optimised to reduce apparent discrepancies between the data products, that would be generated actually by temporal and spatial mismatch and by differences in smoothing of atmospheric variability” are acceptable examples.

4.2.4.4 Information content

A key aspect in the validation of usability (the verification of “fitness for purpose” of a data product) is the characterisation of the information content of the data product. The retrieval of geophysical quantities from remote sounding measurements usually uses a set of a priori constraints, e.g., in the form of an assumed range of atmospheric profile shapes around a first guess. Such constraints mix somehow in the retrieved quantities with the information really contributed by the measurement. When a climatology is used in the retrieval, e.g., at altitudes where the measurement is either not at all or less sensitive due to optically thick clouds or too low signal-to-noise ratios, it is important to understand what information, in the final product, is derived from the climatology and what is really contributed by the measurement. That kind of validation of the information content can rely on a combination of (1) comparisons with independent reference data sets, especially during events not considered in the climatology, (2) the study of deviations of the retrieved product from the a priori constraints, and (3) sensitivity analysis of the retrieval, e.g., based on a study of the associated averaging kernels and their eigenvectors [RD41]. For example, plotting as a function of altitude the sum of the rows of the averaging kernel matrix associated with a retrieval shows at which altitudes the measurement offers sensitivity to atmospheric concentrations. Similarly, the real information content of the reference measurement itself should be known prior to performing a comparison. Information content studies might be an important aspect of the validation of model runs that have been initialised by climatology or by the output of another model, or that are constrained by a priori boundary conditions. They can also be of relevance in the assessment of data assimilation results when observations outside of a predetermined range are rejected as outliers by the data ingestion scheme, producing in the system a zero information zone similar to the dead band or neutral zone used in voltage regulators and controllers to avoid unwanted oscillations and disruptions. Information content studies are also essential in understanding data products generated by data merging and ensemble approaches.



4.2.5 Validation of individual components

ECV product line components are the individual processing blocks by which ECV data products are generated in their interim or final version. For complex processing chains, international standards require researchers to validate or at least verify the good performance of every component and the accuracy of its output. Limiting validation to the final data product only is not sufficient. The validation of intermediate data products is highly desirable to avoid, e.g., that the apparently good behaviour of the final data product at the end of the chain hides large compensating errors affecting separate components of the data retrieval. Testing is one of many verification activities intended to confirm that software development output meets its input requirements. Other verification activities include various static and dynamic analyses, code and document inspections, walkthroughs, and other techniques.

4.2.6 Validation against service specifications

Service specifications are outlined in several documents like the Product Specification Documents (PSD) and the Algorithm Theoretical Basis Documents (ATBD). Verification of every product specification is out of scope of the project. The focus will be on service specifications having clear links with climate research user requirements expressed in the URD [RD5].

4.2.7 Validation against user requirements

User requirements are defined in the URD [RD5], on which summary tables reproduced in Section 3 are based. Products need to be validated against these official user requirements. Assessment of compliance with requirements on observation frequency is straightforward. Compliance with requirements on total uncertainty can be verified by classical comparisons yielding bias and precision estimates, taking into account comparison error terms. Compliance with requirements on spatial resolution and spatial sampling need visualisation of the data and analysis of the information content. Compliance with more specific requirements, e.g., in terms of actual geographical coverage and of point-to-zone representativeness, may require the use of statistical methods based on global model results. In addition to quality checks on the part of the validation teams and the ECV data producers and based on known user requirements, user feedbacks provide valuable input for the assessment of the ECV product compliance in terms of the accuracy (bias, precision or other estimates) and the effective usability of the data product.

4.2.8 Quality control of operational ECV data production

Continuous monitoring of each production line component (e.g., retrieval, modelling, assimilation processes, etc.) within the entire process chain is required (online validation). This comprises monitoring of the operational workflow as well as a permanent quality check of the resulting products. Process failures and data losses have to be documented. Generally, the focus of offline services will be put more on product accuracy, whereas near-real time services (NRT) will be also assessed on the basis of their operational functioning (delay time, loss rate, etc.). In particular, NRT services require access to online available independent measurements from operational networks for automatic validation.

4.2.9 Validation of ECV product updates

Whenever a major upgrade of an ECV data production line occurs (switch to a new sensor, improved retrieval algorithm, updated spectroscopic databases, higher grid resolution...), steps 1 to 3 of the validation in the build-up phase have to be performed and documented: validation of individual components, against service specifications, and against user requirements. The focus must be on the verification of expected product changes. A verification of the entire processing chain might be required as well. A record of successive updates and corresponding validations should be maintained and made publicly available by the ECV data producer. The ECV data producer has to exercise judgement as to the extent of validation needed for a particular service revision, as this will depend on the nature and importance of changes being made. It is also not feasible to test all changes in advance: e.g., sudden degradation of a satellite instrument may necessitate emergency removal of that source from a near-real-time production process.



4.3 Validation specifics by ECV product

4.3.1 Total ozone data product

4.3.1.1 Validation requirements

Validation studies and resulting documentation will address the following targets:

- Time series of ECV total ozone data and of the main measurement and retrieval parameters with potential impact on the data quality (AMF, cloud properties, SZA...) should be visualised, at least in selected latitude zones and at a few representative ground-based stations. Any obvious quality issue like the frequent occurrence of outliers and unrealistic values should be detected, documented and filtered out appropriately before performing quantitative comparisons.
- Statistical estimators of the difference like the bias and the dispersion shall be calculated over different periods and over different ranges of relevant parameters as listed below. In case of frequent occurrence of outliers, median and interpercentile values shall be preferred over mean and standard deviation values as they reduce the influence of outliers. Calculation of mean values and the associated standard deviation is nevertheless encouraged. In case of doubt, histograms of the relative difference might be helpful in determining the validity of statistical estimators.
- In the treatment of statistics, care will be given to decouple as far as possible the different sources of ECV product uncertainty and avoid misleading cancellation of mutually compensating errors. In particular, the dependence of the ECV data quality on main measurement and retrieval parameters like the solar zenith angle, ozone column amount, latitude, and cloud parameters (fractional cloud cover, cloud top height and albedo, etc. as appropriate) shall be investigated.
- Decadal stability of the bias shall be assessed and expressed in %/decade.
- Based on at least bi-weekly sampling of the time series over at least five years, shorter term stability of the bias and dispersion shall be assessed, including annual cycle, interannual variability and shorter-term variability of the bias.
- Studies shall be carried out at least in three geographical zones, in both hemispheres: tropics, middle latitudes and polar areas. Higher meridian and regional sampling is encouraged where possible.

4.3.1.2 Validation data sources

The DARD [RD2] describes the reference measurements to be used for validation studies and/or for cross-comparison studies, with details on their access. The following measurement data sets will be used:

- Ground-based ozone column measurements by Dobson and Brewer UV spectrophotometers, up to 80° SZA for Brewers MK-III and MK-IV and 70-75° of SZA for Dobson instruments and other Brewers.
- Satellite ozone column data by non-CCI retrieval algorithms for EOS-Aura OMI, Metop GOME-2 series, and Suomi-NPP OMPS-NM.



4.3.2 Ozone profile data product from nadir-viewing instruments

4.3.2.1 Validation requirements

Validation studies and resulting documentation will address the following targets:

- Time series of ECV ozone profile data and of the main measurement and retrieval parameters with potential impact on the data quality (SZA, cloud properties...) should be visualised, at least in selected latitude zones and at a few representative ground-based stations. Any obvious quality issue like the frequent occurrence of outliers and unrealistic values should be detected, documented and filtered out appropriately before performing quantitative comparisons.
- Information content issues like the long-term degradation of the Degree of Freedom of the System (DFS) will be studied based on the analysis of vertical averaging kernels and, where relevant, of deviations from the a priori profile.
- Studies shall address ozone in the troposphere, in the UTLS and in the middle atmosphere.
- The error bar on ozone concentration/partial column shall be assessed and expressed as the percent relative difference with respect to correlative measurements of reference. Uncertainties on height registration shall be expressed as the deviation of the retrieval altitude, as expressed by the centroid or the peak altitude of the averaging kernels, from the nominal retrieval altitude. Dependences on time, SZA, latitude, etc. should be identified.
- Statistical estimators of the difference like the bias and the dispersion shall be calculated over different periods and over different ranges of relevant parameters as listed below. In case of frequent occurrence of outliers, median and interpercentile values shall be preferred over mean and standard deviation values as they reduce the influence of outliers. Calculation of mean values and associated standard deviation is nevertheless encouraged. In case of doubt, histograms of the relative difference might be helpful in determining the validity of statistical estimators.
- In the treatment of statistics, care will be given to decouple as far as possible the different sources of ECV product uncertainty and avoid misleading cancellation of mutually compensating errors. In particular, the dependence of the ECV data quality on main measurement and retrieval parameters like the solar zenith angle, ozone slant column amount and latitude shall be investigated.
- Decadal stability of the bias and spread shall be assessed and expressed in %/decade.
- Based on at least bi-weekly sampling of the time series over at least five years, shorter term stability of the bias and dispersion shall be assessed, including annual cycle, interannual variability and shorter term variability of the bias.
- Studies shall be carried out at least in three geographical zones: tropics, middle latitudes and polar areas. Higher meridian and regional sampling is encouraged where possible.

4.3.2.2 Validation data sources

The DARD [RD2] describes the reference measurements to be used for validation studies and/or for cross-comparison studies, with details on their access. The following measurement data sets will be used:

- Ground-based ozone profile measurements by balloon-borne ozonesondes.
- Ground-based ozone profile measurements by stratospheric ozone lidars.



4.3.3 Ozone profile data product from limb-viewing instruments

4.3.3.1 Validation requirements

Validation studies and resulting documentation will address the following targets:

- Time series of ECV ozone profile data and of the main measurement and retrieval parameters with potential impact on the data quality (e.g., SZA for SCIAMACHY) should be visualised, at least in selected latitude zones and at a few representative ground-based stations. Any obvious quality issue like the frequent occurrence of outliers and unrealistic values should be detected, documented and filtered out appropriately before performing quantitative comparisons.
- Studies shall address at least ozone in the lower stratosphere and in the middle atmosphere.
- Statistical estimators of the difference like the bias and the dispersion shall be calculated over different periods and over different ranges of relevant parameters as listed below. In case of frequent occurrence of outliers, median and interpercentile values shall be preferred over mean and standard deviation values as they reduce the influence of outliers. Calculation of mean values and associated standard deviation is nevertheless encouraged. In case of doubt, histograms of the relative difference might be helpful in determining the validity of statistical estimators.
- In the treatment of statistics, care will be given to decouple as far as possible the different sources of ECV product uncertainty and avoid misleading cancellation of mutually compensating errors. In particular, the dependence of the ECV data quality on measurement and retrieval parameters shall be investigated.
- Decadal stability of the bias shall be assessed and expressed in %/decade.
- Based on at least bi-weekly sampling of the time series over at least five years, shorter term stability of the bias and dispersion shall be assessed, including annual cycle, interannual variability and shorter-term variability of the bias.
- Studies shall be carried out at least in three geographical zones: tropics, middle latitudes and polar areas. Higher meridian and regional sampling is encouraged where possible.

4.3.3.2 Validation data sources

The DARD [RD2] describes the reference measurements to be used for validation studies and/or for cross-comparison studies, with details on their access. The following measurement data sets will be used:

- Ground-based ozone profile measurements by balloon-borne ozonesondes.
- Ground-based ozone profile measurements by stratospheric ozone lidars.
- Ground-based ozone profile measurements by ozone microwave radiometers.



4.3.4 Tropospheric ozone data products

4.3.4.1 Validation requirements

Validation studies and resulting documentation will address the following targets:

- Time series of ECV ozone profile data and of the main measurement and retrieval parameters with potential impact on the data quality (SZA, cloud properties...) should be visualised, at least in selected latitude zones and at a few representative ground-based stations. Any obvious quality issue like the frequent occurrence of outliers and unrealistic values should be detected, documented and filtered out appropriately before performing quantitative comparisons.
- Studies shall address ozone in the troposphere as an integrated column.
- The error bar on the tropospheric ozone column shall be assessed and expressed as the percent relative difference with respect to correlative measurements of reference. Dependences on time, SZA, latitude, etc. should be identified.
- Statistical estimators of the difference like the bias and the dispersion shall be calculated over different periods and over different ranges of relevant parameters as listed below. In case of frequent occurrence of outliers, median and interpercentile values shall be preferred over mean and standard deviation values as they reduce the influence of outliers. Calculation of mean values and associated standard deviation is nevertheless encouraged. In case of doubt, histograms of the relative difference might be helpful in determining the validity of statistical estimators.
- In the treatment of statistics, care will be given to decouple as far as possible the different sources of ECV product uncertainty and avoid misleading cancellation of mutually compensating errors. In particular, the dependence of the ECV data quality on main measurement and retrieval parameters like the solar zenith angle and latitude shall be investigated.
- Decadal stability of the bias and spread shall be assessed and expressed in %/decade.
- Based on at least bi-weekly sampling of the time series over at least five years, shorter term stability of the bias and dispersion shall be assessed, including annual cycle, interannual variability and shorter term variability of the bias.
- For low-Earth orbiting (LEO) sounders, studies shall be carried out at least in three geographical zones: tropics, middle latitudes and polar areas. Higher meridian and regional sampling is encouraged where possible. For geostationary (GEO) sounders, studies are spatially limited to the GEO field-of-regard, but tropospheric ozone column observations are available multiple times per day. Comparisons to ozonesondes will at least be made for the closest satellite observation in time, but can be temporally extended to have the daily sampling time as an influence quantity. Daily revisit times will be verified against user requirements.

4.3.4.2 Validation data sources

The DARD [RD2] describes the reference measurements to be used for validation studies and/or for cross-comparison studies, with details on their access. The following measurement data sets will be used:

- Ground-based ozone profile measurements by balloon-borne ozonesondes (vertically integrated up to product-specific tropopause definition).



5 Standards

5.1 *Maintenance of datasets and reports*

It is essential to ensure long-term archiving of ECV data products and their metadata, of validation results and of associated metadata on the validation process, all needed to qualify the stored products and guarantee their proper use in the future and by a widening community. This is achieved by relying on operational archiving systems of the service providers and on the Ozone_cci web site.

I/O documentation and tools for the formats of end products are provided by the ECV data producers. Formats are selected in agreement with the users (netCDF [RD17] in the Ozone_cci project).

5.2 *Metadata and additional information*

Important information on ECV data and their quality must be readily accessible. Beyond comparison results obtained as part of a geophysical validation process, important information covers evaluation from the point of view of the source, technical attributes, quality levels and use conditions, in order to be able to determine whether the data and service are fit for their particular purpose.

Some of this information may be readily available as metadata, but additional information should also be made available if requested to allow an assessment of fitness for purpose to be made. This is particularly important when the data is being used for a purpose, which is different from that for which it was originally produced or collected.

Metadata, whether applying to a dataset or to a service, are necessary for users to:

- Identify and locate the datasets or services they need (“discovery metadata”).
- Be aware of the general context through which the data was collected and made available (research project, programme, etc.), of possible access conditions and of applicable usage rules (such as acknowledgement or citation).
- Retrieve and read the data (format metadata) or access the products provided.
- Understand and interpret the data and their limitations (scientific metadata).
- Seek further information or help if required (references, links, and contact).

In order to fulfil what is expected from them, metadata should be:

- Specific: achieving the level of detail required to an in-depth understanding.
- Accurate: achieving a level of precision sufficient to avoid ambiguities – “accurate” and “precision” here refer to qualities of the wording, not to data.
- Explicit: avoiding coded information, abbreviations and acronyms unless appropriate keys are provided.
- Complete: covering all relevant information, with no omission.

5.3 *QA and validation metadata*

To facilitate proper interpretation of the validation results, traceability of the validation process is essential. Therefore, validation metadata, that is, brief but unambiguous documentation of the entire validation process leading to a validation graph or a comparison data file, should accompany any validation result



reported in validation reports and on the project web site. Where validation results are provided in graphical format (e.g., in a .png file), validation metadata can be provided in a legend placed on the graph itself or below; they can also be attached to the graphical file as a readme.file.text. Where validation results are provided in numerical format (e.g., in an ASCII or HDF file), validation metadata can be included in this numerical data file as a header or simply attached externally to the file.

The metadata on the validation process must provide a short, unambiguous description of the comparison manipulations undertaken to obtain the validation results. From this information, one should be able to check if the validation process complies with agreed standards and best practices. The systematic description of the data manipulations should also allow proper interpretation of the comparison results and further investigation of the data quality.

Table 1 suggests the minimum information that should be available in the validation metadata to ensure traceability of the validation process. Ideally it should not duplicate information that is already available, e.g., in the metadata accompanying the data under evaluation and the validation source.

5.4 *Compliance with international standards*

Interoperability is a driving concept of the GEOSS Implementation Plan in general and of the CCI/CCI+ programme in particular. Elaborated in this context, the present document gives particular attention to international standardisation requirements formulated e.g. within high-level strategies like the QA4EO framework formalised by the Committee on Earth Observation Satellites (CEOS) and the Integrated Global Observation Strategy (IGOS) established by a list of international partners (including CEOS, GAW, GCOS, IGBP, UNEP, UNESCO, WCRP and WMO), and within European initiatives relevant to GMES. Further evolution is anticipated.

Particular attention must be paid to the Directive 2007/2/EC of the European Parliament and of the Council of 14 March 2007, establishing an Infrastructure for Spatial Information in the European Community (INSPIRE) to support Community environmental policies, and policies or activities which may have an impact on the environment. Published in the Official Journal on the 25th of April 2007, the INSPIRE Directive entered into force on the 15th of May 2007. To ensure that the spatial data infrastructures of the Member States are compatible and usable in a Community and transboundary context, the Directive requires that common Implementing Rules (IR) are adopted in a number of specific areas (Metadata, Data Specifications, Network Services, Data and Service Sharing and Monitoring and Reporting). These IRs are adopted as Commission Decisions or Regulations, and are binding in their entirety. Data themes under Annex III of the IRs have the more direct application to the Ozone_cci+ project, among them: Atmospheric conditions, Environmental monitoring facilities, Statistical units, Human health and safety, Natural risk zones, Meteorological geographic features. The NetCDF formats adopted in Ozone_cci are compliant with INSPIRE IRs.



Table 1. Suggested validation metadata

VALIDATION STEP / ITEM	DETAILS
High-level description of the content of validation results (graphic file or numerical data file)	Identification of the data being validated and of the reference data used as a validation source, date, basic description of the results being reported
Metadata on data under evaluation	Data processing and archiving centre, model or data processor version, input and initialisation data, native data format (e.g., number density or volume mixing ratio, versus altitude or pressure...), data file name (at least file name convention)
Metadata on reference data used as a validation source	Data processing and archiving centre, instrument, responsible institute, model/data processor version, calibration version (input level-1 data), measured parameter, native data format...
Traceability of validation process	Systematic description of the data manipulations: data selection, conversion of units, filtering based, e.g., on flags or statistical tests, co-location criteria (vertical, horizontal and temporal), regridding and smoothing (vertical and horizontal, e.g. using a Gaussian, averaging kernels etc.), domain of the comparisons (geographical, vertical, temporal), reference to an agreed reference practice...
Format of validation results	Content of the numerical validation data file or description of the information displayed on the validation graph: units, relative or absolute difference, individual comparison pair or monthly mean, amount of comparison events, statistical estimators (mean/deviation or median/interpercentile) ...
Credit and responsibilities	Analysis carried out at institute X by validation scientist Y supported by data processing scientist Z, contact (email)...



6 Terms and definitions

6.1 Terms and definitions

In Table 2, terms and definitions as recommended by CEOS WGCV and by standards development organisations of international recognition have been transcribed from reference documents [RD7] to [RD15]. In some cases, terms and definitions peculiar to forecast systems are also proposed. They are expected to evolve as these organisations regularly update their standards and as further standardisation and harmonisation occur.

Table 2. Recommended terms and definitions.

TERM	DEFINITION	SOURCE
accuracy	closeness of agreement between a quantity value obtained by measurement and the true value of the measurand; note that <u>it is not a quantity</u> and it is not given a numerical quantity value	VIM, GUM
area (volume) of representativeness	the area (volume) in which the concentration does not differ from the concentration at the station by more than a specific range	Larssen
bias	(1) systematic error of indication of a measuring system (2) estimate of a systematic measurement error (3) estimate of a systematic forecast error	VIM VIM GAS
calibration	(1) the process of quantitatively defining the system responses to known, controlled signal inputs (2) operation that, under specified conditions, in a first step, establishes a relation between the quantity values with measurement uncertainties provided by measurement standards and corresponding indications with associated measurement uncertainties and, in a second step, uses this information to establish a relation for obtaining a measurement result from an indication	CEOS VIM
dead band (or neutral zone)	maximum interval through which a value of a quantity being measured can be changed in both directions without producing a detectable change in the corresponding indication	VIM
detection limit	measured quantity value, obtained by a given measurement procedure, for which the probability of falsely claiming the absence of a component, given a probability α of falsely claiming its presence	VIM
error	(1) measured quantity value minus a reference quantity value (2) difference of quantity value obtained by measurement and true value of the measurand (3) difference of forecast value and a, estimate of the true value	VIM CEOS
establish	define, document and implement	CDRH
field-of-regard	an area of the object space scanned by the field-of-view of a scanning sensor	NIST
field-of-view	the solid angle from which the detector receives radiation	NIST
footprint	the area of a target encircled by the field-of-view of a detector of radiation, or irradiated by an active system	NIST
influence quantity	quantity that, in a direct measurement, does not affect the quantity that is actually measured, but affects the relation between the indication and the measurement result	VIM



TERM	DEFINITION	SOURCE
in situ measurement	(1) a direct measurement of the measurand in its original place (2) any sub-orbital measurement of the measurand	GEOSS
measurand	quantity intended to be measured	VIM
metadata	data about the data; parameters that describe, characterise, and/or index the data	WMO
monitoring	(1) systematic evaluation over time of some quantity (2) by extension, evaluation over time of the performance of a system, of the occurrence of an event etc.	NIST
point-to-area (point-to-volume) representativeness	the probability that a point measurement lies within a specific range of area-average (volume-average) concentration value	Nappo
precision	closeness of agreement between quantity values obtained by replicate measurements of a quantity on the same or similar object under specified conditions	VIM
process validation	establishing documented evidence of a high degree of assurance that a specific process will consistently produce a product meeting its pre-determined specifications and quality characteristics	CDRH
quality assessment (QA)	QA refers to the overall management of the processes involved in obtaining the data	CEOS
quality control (QC)	QC refers to the activities undertaken to check and optimise accuracy and precision of the data after its collection	CEOS
quality indicator (QI)	a means of providing a user of data or derived product with sufficient information to assess its suitability for a particular application. This information should be based on a quantitative assessment of its traceability to an agreed reference or measurement standard (ideally SI), but can be presented as a numeric or a text descriptor, provided the quantitative linkage is defined.	QA4EO
radiometric calibration	a determination of radiometric instrument performance in the spatial, spectral, and temporal domains in a series of measurements, in which its output is related to the true value of the measured radiometric quantity	NIST
random error	(1) component of measurement error that in replicate measurements varies in an unpredictable manner; note that random measurement error equals measurement error minus systematic measurement error (2) component of forecast error that varies in an unpredictable manner	VIM
relative standard uncertainty	standard measurement uncertainty divided by the absolute value of the measured quantity value	VIM
repeatability	measurement precision under set of conditions including the same measurement procedure, same operator, same measuring system, same operating conditions and same location, and replicated measurements over a short period of time	VIM
representativeness	the extent to which a set of measurements taken in a given space-time domain reflect the actual conditions in the same or different space-time domain taken on a scale appropriate for a specific application	Nappo
reproducibility	measurement precision under a set of conditions including different locations, operators, and measuring systems	VIM



TERM	DEFINITION	SOURCE
resolution	(1) the least angular/linear/temporal/spectral distance between two identical point sources of radiation that can be distinguished according to a given criterion (2) the least vertical/geographical/temporal distance between two identical atmospheric features that can be distinguished in a gridded numerical product or in time series of measurements; resolution is equal to or coarser than vertical/geographical/temporal sampling of the grid or the measurement time series	NIST
stability	ability of a measuring system to maintain its metrological characteristics constant with time	VIM
systematic error	component of measurement error that in replicate measurements remains constant or varies in a predictable manner	VIM
traceability	property of a measurement result relating the result to a stated metrological reference (free definition and not necessarily SI) through an unbroken chain of calibrations of a measuring system or comparisons, each contributing to the stated measurement uncertainty	VIM
tropopause	the region of the atmosphere where the environmental temperature lapse rate changes from positive (in the troposphere) to negative (in the stratosphere) the lowest level at which the lapse rate decreases to 2 °C/km or less, provided that the average lapse rate between this level and all higher levels within 2 km does not exceed 2 °C/km occasionally, a second tropopause may be found if the lapse rate above the first tropopause exceeds 3 °C/km	WMO
uncertainty	non-negative parameter that characterizes the dispersion of the quantity values that are being attributed to a measurand, based on the information used	VIM
validation	(1) the process of assessing, by independent means, the quality of the data products derived from the system outputs (2) verification where the specified requirements are adequate for an intended use (3) the process of assessing, by independent means, the degree of correspondence between the value of the radiometric quantity derived from the output signal of a calibrated radiometric device and the actual value of this quantity. (4) confirmation by examination and provision of objective evidence that specifications conform to user needs and intended uses, and that the particular requirements implemented through software can be consistently fulfilled	CEOS VIM NIST CDRH
verification	(1) the provision of objective evidence that a given data product fulfils specified requirements; note that, when applicable, measurement uncertainty should be taken into consideration. (2) the provision of objective evidence that the design outputs of a particular phase of the software development life cycle meet all of the specified requirements for that phase	VIM CDRH
vicarious calibration	a post-launch radiometric calibration of sensors performed with the use of natural or artificial sites or objects on the surface of the Earth (as opposed to calibration techniques using onboard standards such as lamps, blackbodies, solar diffuse reflecting panels etc.)	NIST



6.2 Abbreviations and acronyms

Note of best practice: Using an acronym is acceptable if it has been defined the first time it appears in a document. The same applies to chemical abbreviations. In documents targeting a wide spectrum of potential readers, like user manuals and validation reports, it is recommended to avoid systematic use of acronyms and abbreviations except for those with frequent occurrence, and those widely understood by the general public. For example, acronyms such as CFCs and ESA are acceptable. Acronyms such as ECSS and ICTT-QMF are not. Before using acronyms and abbreviations, authors should keep in mind that it is annoying and difficult – especially in Web-based documents unless the acronyms are available as hyperlinks – to turn over several pages in a document to verify the meaning.

AK	Averaging Kernel
ALGT	Algorithm development Team
AMF	Air Mass Factor, or optical enhancement factor
ATBD	Algorithm Theoretical Basis Document
AUTH	Aristotle University of Thessaloniki
BIPM	Bureau International des Poids et Mesures
BIRA-IASB	Belgian Institute for Space Aeronomy
CCI	ESA's Climate Change Initiative programme
CDRH	Center for Devices and Radiological Health
CEOS	Committee on Earth Observation Satellites
CMUG	Climate Modelling User Group of the CCI programme
CRG	Climate Research Group of the Ozone_cci project
DARD	Data Access Requirement Document
DFS	Degree of Freedom of the System
DHF	Data Host Facility
DLR	German Aerospace Centre
DOAS	Differential Absorption Optical Spectroscopy
DU	Dobson Unit – unit of vertical column density ($2.69 \cdot 10^{16} \text{ molec.cm}^{-2}$)
EC	European Commission
ECSS	European Corporation for Space Standardization
Envisat	ESA's Environmental Satellite, launched March 1, 2002
EO	Earth Observation
ERS-2	ESA's Earth Remote Sensing satellite 2, launched April 21, 1995
ESA	European Space Agency
ESRIN	European Space Research Institute
EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
FMI	Finnish Meteorological Institute
GAW	WMO's Global Atmosphere Watch
GCOS	Global Climate Observing System
GDP	GOME Data Processor
GEO	Group on Earth Observation
GEOSS	Global Earth Observation System of Systems
GMES	Global Monitoring for Environment and Security
GOME	Global Ozone Monitoring Experiment
GOMOS	Global Ozone Monitoring by Occultation of Stars
GSE	GMES Service Element
GUM	Guide to the expression of uncertainty in a measurement
HALOE	Halogen Occultation Experiment
ICTT-QMF	Inter-Commission Task Team on Quality Management Framework
IGACO	Integrated Global Atmospheric Chemistry Observation strategy
IGBP	International Geosphere-Biosphere Project
IGOS	Integrated Global Observation Strategy



INSPIRE	Infrastructure for Spatial Information in the European Community
I/O	Input/Output
IR	INSPIRE Implementation Rule
ISO	International Organization for Standardization
JCGM	Joint Committee for Guides in Metrology
KNMI	Royal Dutch Meteorological Institute
Lidar	Light detection and ranging
MetOp	EUMETSAT's Meteorological Operational satellite
MIPAS	Michelson Interferometer for Passive Atmospheric Sounding
MLS	Microwave Limb Sounder
MPC	Mission Performance Centre
Multi-TASTE	Technical ASsistance To the multi-mission validation of Envisat and Third Party Missions using spectrometers, radiometers and sondes
NDACC	Network for the Detection of Atmospheric Composition Change
NDSC	Network for the Detection of Stratospheric Change (now NDACC)
NOAA	National Oceanic and Atmospheric Administration
NRT	Near-real time
O ₃	ozone
OMI	Ozone Monitoring Instrument
OPERA	Ozone Profile Retrieval Algorithm
PROMOTE	Protocol Monitoring for the GMES Service Element - Atmosphere
PSD	Product Specification Document
PVP	Product Validation Plan
QA4EO	Quality Assurance framework for Earth Observation
RAL	Rutherford Appleton Laboratory
SAGE	Stratospheric Aerosol and Gas Experiment
SBUV	Solar Backscatter Ultraviolet
SCIAMACHY	SCanning Imaging Absorption spectroMeter for Atmospheric CHartographY
SHADOZ	Southern Hemisphere ADDitional Ozonesondes
SZA	Solar Zenith Angle
TOC	Total Ozone Column
TOMS	Total Ozone Mapping Spectrometer
UARS	Upper Atmosphere Research Satellite, launched September 15, 1991
UNEP	United Nations Environment Programme
UNESCO	United Nations Educational Scientific and Cultural Organization
URD	User Requirement Document
USM	Upper Stratosphere/Mesosphere
UT	Upper Troposphere
UTLS	Upper Troposphere/Lower Stratosphere
VALT	Validation team of the Ozone_cci project
VIM	International Vocabulary of Metrology — Basic and general concepts and associated terms
WCRP	World Climate Research Project
WGCV	CEOS Working Group on Calibration and Validation
WMO	World Meteorological Organization
WOUDC	World Ozone and Ultraviolet Radiation Data Center